

Correction to “Recent Cooling of the Upper Ocean”

Josh K. Willis¹, John M. Lyman^{2,3}, Gregory C. Johnson² and John
Gilson⁴

Submitted 9 April 2007

to

Geophysical Research Letters

Abstract. The recent cooling signal in the upper ocean reported by *Lyman et al.* [2006] is shown to be an artifact that was caused by a large cold bias discovered in a small fraction of Argo floats as well as a smaller but more prevalent warm bias in expendable BathyThermograph (XBT) data. These biases are both substantially larger than sampling errors estimated in *Lyman et al.* [2006].

Index Terms. 4513 Decadal ocean variability; 1225 Global change from geodesy; 4215 Climate and interannual variability; 4262 Ocean observing systems; 1635 Oceans
Keywords. global warming; heat content; climate change

¹ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

² NOAA/Pacific Marine Environmental Laboratory, 7600 Sand Point Way Bldg. 3,
Seattle, Washington 98115-6349, U.S.A.

³ JIMAR, University of Hawaii, Honolulu, HI

⁴ Scripps Institution of Oceanography, University of California, San Diego, La
Jolla, California

The rapid decrease in globally integrated upper (0–750 m) ocean heat content anomalies (OHCA) between 2003 and 2005 reported by *Lyman et al.* [2006] appears to be an artifact resulting from the combination of two different instrument biases recently discovered in the in situ profile data. Although *Lyman et al.* [2006] carefully estimated sampling errors, potential biases between different instrument types were not investigated. One such bias has been identified in a subset of Argo float profiles that can be easily excluded from the OHCA estimates. The other bias, however, appears to be caused by expendable BathyThermograph (XBT) data that are systematically warm [*Gouretski and Koltermann, 2007*] compared to other instruments. Characterization and removal of this bias will be required before the historical record of ocean heat content can be reevaluated.

The first bias is due to a small subset of Argo float profiles for which incorrect pressure values have been assigned. This problem was detected by comparison of individual Argo temperature profiles with historical data in regions of the Atlantic Ocean that appeared to exhibit rapid cooling. Further investigation uncovered a design flaw in a specific type of Argo float. The floats are SOLO (Sounding Oceanographic Lagrangian Observer) instruments fabricated at WHOI (the Woods Hole Oceanographic Institution) and equipped with FSI (Falmouth Scientific, Inc.) CTD (Conductivity-Temperature-Depth) sensors. These floats contain an error in their internal software that causes them to associate temperature and salinity with pressure values that tend to be biased shallow, but not uniformly so over the entire profile. This error results in profiles that are spuriously cold, with the bias being largest around the depth of the thermocline. The bias affects approximately 14,000 float profiles (about 6 % of all Argo profiles) from 287 individual floats deployed primarily in the Atlantic Ocean between mid-2003 and 2006.

Using additional information reported by the float, approximately 30 % of the biased profiles will be corrected exactly, and the remainder will be corrected approximately (B. Owens, personal communication, 2007). As of this writing, data have not yet been corrected and must therefore be excluded from the OHCA analysis. The floats can be identified in two different ways. First, these floats comprise the dominant subset of WMO (World Meteorological Organization) instrument type 852 in the Argo data set. Second, the individual WMO identification numbers of the floats have been published on the internet (see <http://www.argo.ucsd.edu/> for these numbers and more details).

The second instrument bias arises in data from expendable BathyThermograph (XBT) instruments. These inexpensive probes were not designed to provide climate-quality scientific data. Nevertheless, data from these instruments make up a large fraction of historical ocean temperature measurements, and efforts have been made to remove systematic biases such as those caused by errors in the equations used to model XBT fall rates [*Hanawa et al., 1995*]. Recent work by *Gouretski and Koltermann* [2007], however, suggests that a warm bias remains in the XBT data set. After removing the bias caused by the spurious float data, we find evidence for a warm bias in the XBT data as well.

The effects of these biases are evident in estimates of the globally integrated OHCA made from subsets of the profile data (Figure 1). These estimates were computed by integrating over maps of OHCA made using the techniques and mapping parameters described by *Willis et al.* [2004]. Seasonal temperature anomalies from the World Ocean

Atlas 2001 [Stephens *et al.*, 2001] and a 1993–2003 climatology were used as the first guess. As in Lyman *et al.* [2006], the error bars represent one standard error based on sampling error only. Additional uncertainties may arise from instrument biases and errors in the climatological mean and seasonal cycle.

The OCHA estimate made using all data (thick solid line), including spurious float profiles, shows an apparent cooling of about 48×10^{21} J (48 zetajoules) from 2004 through 2006. This erroneous cooling arises because of the increasing fraction of spurious profiles in the Argo data stream produced by the WHOI SOLO floats. Another estimate using all data except the spurious float data (thick dashed line) suggests much less cooling, only about 25 zetajoules.

The effect of the XBT bias is demonstrated by making OHCA estimates from two more subsets of the data. One is made by excluding all Argo float data (thin dashed line), and consists primarily of XBT profiles. The other is made using only Argo data but excluding the spurious float profiles (thin solid line). The start year of 2004 was chosen because it was the first year in which Argo floats achieved adequate coverage to compute the global integral of OHCA. Although neither estimate shows significant warming or cooling during this period, they are separated by 76 zetajoules, much greater than the sampling error in either estimate. Here all OHCA estimates are plotted relative to the same mean, whereas in Lyman *et al.* [2006; their Fig. 1], record-length means differing by 14 zetajoules were subtracted from the two different OHCA estimates before plotting. The significant separation between the XBT and Argo based estimates supports the existence of a warm bias in the XBT data as reported by Gouretski and Koltermann [2007]. The reason for the apparent cooling in the estimate that combines both of these data sets (thick dashed line) is the increasing ratio of Argo observations to XBT observations between 2004 and 2006. The combined estimate therefore exhibits cooling as it moves away from the warm XBT data and toward the cooler Argo values.

Unlike the Argo pressure problem, the cause of the warm bias in the XBT data remains uncertain. Furthermore, work by Gouretski and Koltermann [2007] suggests that it may vary in both time and depth making it difficult to remove. Despite its shortcomings, the XBT network provides an important means of monitoring basin-wide mass and heat transports in the ocean [Douglas *et al.*, 2006; Gilson *et al.*, 1998] and has dramatically improved sampling of globally integrated OHCA since the early 1970s [Lyman *et al.*, 2006]. Careful calibration of the XBT data against other data types is therefore needed in order to reduce the systematic biases that remain.

However, until the XBT warm bias is addressed, combining XBT data with data from the Argo array to produce annual estimates of OHCA is not prudent. Lack of any significant cooling signal in either the Argo-based or XBT-based estimates suggests that the cooling reported by Lyman *et al.* [2006] was the result of both the cold bias in the spurious Argo data as well as the transition from OHCA estimates dominated by warm XBT data to estimates dominated by Argo data. Finally, removal of the upper-ocean cooling signal also brings estimates of thermosteric sea level variability into closer agreement with altimeter-derived measurements of global mean sea level rise, requiring much less dramatic increases in the melting of land bound ice to close the sea level budget.

Discovery of these biases highlights the need for careful intercalibration of data streams in the global ocean observing system. As with all climate research, maintaining

the capabilities of the ocean observing system requires that significant resources be devoted not only to maintaining high quality observing systems such as the Argo array, but also to the intercomparison and analysis of different data streams as observing systems evolve.

Acknowledgments. The bulk of the in situ data used herein were provided through the World Ocean Database 2001 and the Global Temperature-Salinity Profile Program (<http://www.nodc.noaa.gov>). Float data were collected and made freely available by Argo (a pilot program of the Global Ocean Observing System) and contributing national programs (<http://www.argo.net/>). JML and GCJ were supported by the NOAA Climate Program Office and the NOAA Office of Oceanic and Atmospheric Research. The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the National Oceanic and Atmospheric Administration. This research was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. PMEL contribution 3069. JIMAR Contribution 07-364.

Reference List

- Douglass, E., D. Roemmich, and D. Stammer (2006), Interannual variability in northeast Pacific circulation, *J. Geophys. Res.*, *111*, C04001, doi:10.1029/2005JC003015.
- Gilson, J., D. Roemmich, B. Cornuelle, and L.-L. Fu (1998), Relationship of TOPEX/Poseidon altimetric height to steric height and circulation of the North Pacific, *J. Geophys. Res.*, *103*, 27,947–27,965.
- Gouretski V., K. P. Koltermann (2007), How much is the ocean really warming?, *Geophys. Res. Lett.*, *34*, L01610, doi:10.1029/2006GL027834.
- Hanawa, K., P. Rual, R. Bailey, A. Sy, and M. Szabados (1995), A new depth-time equation for Sippican or TSK T-7, T-6, and T-4 expendable bathythermographs (XBT), *Deep-Sea Res. I*, *42*, 1423–1451.
- Lyman, J. M., J. K. Willis, G. C. Johnson (2006), Recent Cooling in the Upper Ocean, *Geophys. Res. Lett.*, *33*, L18604, doi:10.1029/2006GL027033.
- Stephens, C., J. I. Antonov, T. P. Boyer, M. E. Conkright, R. A. Locarnini, T. D. O'Brien, and H. E. Garcia (2002), World ocean atlas 2001, volume 1: Temperature, *Tech. Rep., Atlas NESDIS 49*, 167 pp., Natl. Ocean and Atmos. Admin., Washington, D.C.
- Willis, J. K., D. Roemmich, and B. Cornuelle (2004), Interannual variability in upper ocean heat content, temperature, and thermosteric expansion on global scales, *J. Geophys. Res.*, *109*, C12036, doi:10.1029/2003JC002260.

Figures




Figure 1. Annual values of globally averaged OHCA in the upper 750 m using all available data (thick solid line), OHCA using all data except profiles from SOLO floats with FSI sensors (thick dashed line), OHCA using only Argo data except SOLO floats with FSI sensors (thin solid line), OHCA with no Argo data (thin dashed line). As in Lyman *et al.* [2006], error bars reflect only sampling errors and not the complete error budget.

Recent Cooling of the Upper Ocean

John M. Lyman^{1,2,3}, Josh K. Willis⁴, and Gregory C. Johnson¹

Submitted 26 May 2006

to

Geophysical Research Letters

Accepted 31 July 2006

Published 20 September 2006

Abstract. We observe a net loss of $3.2 (\pm 1.1) \times 10^{22}$ J of heat from the upper ocean between 2003 and 2005. Using a broad array of in situ ocean measurements, we present annual estimates of global upper-ocean heat content anomaly from 1993 through 2005. Including the recent downturn, the average warming rate for the entire 13-year period is 0.33 ± 0.23 W/m² (of the Earth's total surface area). A new estimate of sampling error in the heat content record suggests that both the recent and previous global cooling events are significant and unlikely to be artifacts of inadequate ocean sampling.

Index Terms. 4513 Decadal ocean variability; 1225 Global change from geodesy; 4215 Climate and interannual variability; 4262 Ocean observing systems; 1635 Oceans
Keywords. global warming; heat content; climate change

¹ NOAA/Pacific Marine Environmental Laboratory, 7600 Sand Point Way N.E. Bldg. 3, Seattle Washington 98115-6349, U.S.A.

² JIMAR, University of Hawaii, Honolulu, HI

³ Corresponding author. Tel.: +1-206-526-4420; fax: +1-206-526-6744, E-mail address: John.Lyman@noaa.gov (J. M. Lyman).

⁴ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA

1. Introduction

With over 1000 times the heat capacity of the atmosphere, the World Ocean is the largest repository for changes in global heat content [Levitus *et al.*, 2005]. Monitoring ocean heat content is therefore fundamental to detecting and understanding changes in the Earth's heat balance. Past estimates of the global integral of ocean heat content anomaly (OHCA) indicate an increase of 14.5×10^{22} J from 1955 to 1998 from the surface to 3000 m [Levitus *et al.*, 2005] and $9.2 (\pm 1.3) \times 10^{22}$ J from 1993 to 2003 in the upper (0 – 750 m) ocean [Willis *et al.* 2004]. These increases provide strong evidence of global warming. Climate models exhibit similar rates of ocean warming, but only when forced by anthropogenic influences [Gregory *et al.*, 2004; Barnett *et al.*, 2005; Church *et al.*, 2005; Hansen *et al.*, 2005].

While there has been a general increase in the global integral of OHCA during the last half century, there have also been substantial decadal fluctuations, including a short period of rapid cooling (6×10^{22} J of heat lost in the 0–700 m layer) from 1980 to 1983 [Levitus *et al.*, 2005]. Most climate models, however, do not contain unforced decadal variability of this magnitude [Gregory *et al.*, 2004; Barnett *et al.*, 2005, their Figure S1; Church *et al.*, 2005; and Hansen *et al.*, 2005] and it has been suggested that such fluctuations in the observational record may be due to inadequate sampling of ocean temperatures [Gregory *et al.*, 2004]. We have detected a new cooling event that began in 2003 and is comparable in magnitude to the one in the early 1980s. Using high-resolution satellite data to estimate sampling error, we find that both the recent event and the cooling of the early 1980s are significant with respect to these errors.

2. Heat Content Anomaly

Using a broad array of in situ temperature data from expendable bathythermographs (XBTs), ship board conductivity-temperature-depth (CTD) sensors, moored buoy thermistor records (primarily from Tropical Atmosphere Ocean array), and autonomous profiling CTD floats (primarily from Argo) the global integral of OHCA of the upper 750 m is estimated from the start of 1993 through the end of 2005 (Figure 1). The global integral is computed from 1/4 degree mapped fields of annual averaged OHCA as in Willis *et al.* [2004], except that in the present analysis the altimeter data are not used. From 1993 to 2003, the heat content of the upper ocean increased by $8.1 (\pm 1.4) \times 10^{22}$ J. This increase was followed by a decrease of $3.2 (\pm 1.1) \times 10^{22}$ J between 2003 and 2005. The decrease represents a substantial loss of heat over a 2-year period, amounting to about one fifth of the long-term upper-ocean heat gain between 1955 and 2003 reported by Levitus *et al.*, [2005].

From 1993 to 2005, the average rate of upper-ocean warming as determined by a linear least squares fit is 0.33 ± 0.23 W/m² (of the Earth's total surface area). This convention is chosen to emphasize the observationally supported relationship between ocean heat content and the Earth's energy balance [Pielke, 2003; Levitus *et al.*, 2005; Wong *et al.*, 2006]. The uncertainty represents the 95% confidence interval and reflects both the random error in each annual estimate as well as the interannual variability in the curve that is not explained by a linear trend. To calculate the uncertainty, the effective degrees of freedom were computed by dividing the length of the time series by the

decorrelation length scale of the residuals from the fit. The recent decrease in heat content amounts to an average cooling rate of $-1.0 \pm 0.3 \text{ W/m}^2$ (of the Earth's total surface area) from 2003 to 2005, and results in a lower estimate of average warming from 1993 to 2005 than that recently reported for the 1993 to 2003 period [Willis *et al.*, 2004]. It is important to note that this decrease causes greater uncertainty in the long-term warming rate because the cooling reflects interannual variability that is not well represented by a linear trend. This cooling event, as well as the cooling in the early 1980s, illustrates the importance of accounting for interannual variability when determining long-term rates of ocean warming. This interannual variability complicates the task of detecting upper ocean warming due to anthropogenic influence, which is assumed to have a time scale of many decades.

The recent downturn in OHCA roughly coincides with the spin up of Argo [*www.argo.net*] in 2002. Argo has dramatically improved sampling and introduced a large amount of data from new instruments, namely autonomous profiling CTD floats. In order to test for potential biases due to this change in the observing system, globally averaged OHCA was also computed without profiling float data (Figure 1, gray line). The cooling event persisted with removal of all Argo data from the OHCA estimate, albeit more weakly and with much larger error bars. This result suggests that the cooling event is real and not related to any potential bias introduced by the large changes in the characteristics of the ocean observing system during the advent of the Argo Project. Estimates of OHCA made using only data from profiling floats (not shown) also yielded a recent cooling of similar magnitude.

The relatively small magnitude of the globally averaged signal is dwarfed by much larger regional variations in OHCA (Figure 2). These variations sometimes exceed the equivalent of a local air-sea heat flux anomaly of 50 W/m^2 applied continuously over 2 years and so are too large to be caused by this mechanism alone. Changes such as these are also due to mesoscale eddy advection, advection of heat by large-scale currents, and interannual to decadal shifts in gyre circulation that are associated with climate phenomena such as El Niño [Johnson *et al.*, 2000], the North Atlantic Oscillation [Curry *et al.*, 2001], the Pacific Decadal Oscillation [Deser *et al.*, 1999], and the Antarctic Oscillation [Roemmich *et al.*, 2006]. Owing in part to the strength of these advection-driven changes, the source of the recent globally averaged cooling (Figure 1) cannot be localized from OHCA data alone.

3. Uncertainty in the Global Integral

Assessing the significance of the comparatively tiny (order 1 W/m^2) changes in the global average OHCA requires an estimate of how well the large regional signals are resolved by the often sparsely sampled in situ OHCA data. Since late 1992, dense, high-quality measurements of sea surface height anomaly (SSHA) have been obtained via satellite altimeter. Maps of SSHA from Aviso (a combined satellite altimeter product) contain variability on scales as small as ten days and 150 – 200 km [Ducet *et al.*, 2000], have almost complete global coverage (excluding ice-covered regions), and are related to ocean heat content [White *et al.*, 1996; Gilson *et al.*, 1998; Willis *et al.*, 2004]. Admittedly, SSHA variability is not perfectly correlated with OHCA variability. SSHA variations are also influenced by ocean freshwater content owing to precipitation, evaporation, and run-off, as well as by deep ocean variations (below 750 m in this case). Despite these complicating factors, the correlation between SSHA and OHCA holds

reasonably well and is used here to compute the uncertainty in the in situ estimate of OHCA.

Estimates of the uncertainty in OHCA are made for the years 1955 to 2005 by sub-sampling the 13-year record of SSHA in the same manner as the in situ sampling pattern for a given year, N . The global integral of SSHA, for the 13-year record, is constructed from maps made from the sub-sampled data set [Willis *et al.*, 2004] and compared to the global integral of SSHA based on the complete maps of Aviso data. Taking the time series based on the complete maps as truth, the uncertainty for year N is expressed as a standard deviation:

$$sampling_error(N) = 5.1 \times 10^{22} J \cdot cm^{-1} \left[\frac{\sum_{i=1993}^{2005} [SSH_{total}(i) - SSH_{sub_N}(i)]^2}{13} \right]^{1/2} \quad (1)$$

where the proportionality constant between SSHA and OHCA is $5.1 \times 10^{22} J \cdot cm^{-1}$ [Willis *et al.*, 2004], SSH_{total} is the global average of SSHA from the complete maps for year i , and SSH_{sub_N} is the global average of SSHA from the Aviso data for year i sub-sampled at observation locations for year N and then remapped. There is only one realization of the globally averaged OHCA each year, therefore the standard deviation of the sampling error and the standard error are the same.

This method most likely underestimates the sampling error, as the 13-year record of SSHA is missing variability from time scales longer than decadal and shorter than 10 days. The method also assumes that the shorter time scale variability in the decades preceding the 1990s is similar to that from 1993 to 2005, which may not be true. Hence, the error estimate is probably most accurate for the period of satellite altimetry, since 1993. Despite these caveats, this process likely produces a reasonable estimate of the sampling error in one-year averages of OHCA prior to 1993 as well. It is worth noting, however, that lack of a longer altimeter record may preclude using this technique to determine accurate uncertainties for the long-term warming rate reported by Levitus *et al.* [2005] that has been the subject of recent debate [Gregory *et al.*, 2004; Gille, 2006].

The standard error on the annually averaged global mean from Aviso SSH maps [Willis *et al.*, 2004] is $0.2 \times 10^{22} J$. This term is combined with the standard error from the sampling error computed above, assuming these two errors are independent, to yield the standard error on the OHCA estimate (Figure 3) for a given year N :

$$standard_error_OHCA(N) = \left[(0.2 \times 10^{22} J)^2 + sampling_error(N)^2 \right]^{1/2}. \quad (2)$$

The time-period from 1955 to 2005 can be broken into three different epochs with regards to in situ sampling of OHCA. The first epoch, prior to the advent of XBTs, ended around 1967. Globally averaged uncertainty during this epoch (Figure 3) is on the same order as the decadal signal [Levitus *et al.*, 2005] making it difficult to quantify decadal changes in the globally averaged OCHA prior to 1968.

Upon the commencement of widespread use of XBTs in 1968, a second epoch began that continued until 2002. Uncertainty in globally averaged OHCA drops by a factor of six from 1955 to 1968 (Figure 3). The decrease in uncertainty is due to the increase in the number of observations from 4,500 in 1955 to 31,900 in 1968, an increase that was fueled by the introduction of the XBT. During the second epoch, the error

decreases only slightly with time (Figure 3) but is generally small compared to decadal changes in globally averaged OHCA. In particular, the 6×10^{22} J decrease in heat content during the early 1980s that was reported by *Levitus et al.* [2005] lies well outside the range of uncertainty presented here.

A third epoch began around 2003 with the ramp-up of Argo. The goal of this international project is to deploy and maintain an array of 3000 autonomous profiling CTD floats designed to accurately measure temperature and salinity in the upper 2000 m of the global ice-free ocean at 10-day intervals and $3^\circ \times 3^\circ$ spatial resolution. From 2002 to 2005 there was a factor of three decrease in the standard error of OHCA that resulted directly from Argo data. The uncertainty in the global average of annual OHCA is now at a historic low of 0.6×10^{22} J. Thus, the magnitude of the recent cooling is also well outside the range of uncertainty. While the number of in situ samples is about the same in 2002 and 2005, the latter year is sampled by well-dispersed Argo floats and has a much more even distribution of data compared to 2002 (Figure S1). In addition, because Argo floats report back in real time, near real-time estimates of OHCA are now possible.

4. Vertical Structure of Cooling

The depth structure of globally averaged temperature change between 2003 and 2005 (Figure 4) allows a few more insights into the recent cooling. Uncertainty in the temperature change was computed by scaling the 1×10^{22} J standard error in the heat content decrease using regression coefficients for temperature variability [*Willis et al.*, 2004]. The average uncertainty is about 0.01 °C at a given depth. The cooling signal is distributed over the water column with most depths experiencing some cooling. A small amount of cooling is observed at the surface, although much less than the cooling at depth. This result of surface cooling from 2003 to 2005 is consistent with global SST products [e.g. http://www.jisao.washington.edu/data_sets/global_sstanomts/]. The maximum cooling occurs at about 400 m and substantial cooling is still observed at 750 m. This pattern reflects the complicated superposition of regional warming and cooling patterns with different depth dependence, as well as the influence of ocean circulation changes and the associated heave of the thermocline.

The cooling signal is still strong at 750 m and appears to extend deeper (Figure 4). Indeed, preliminary estimates of 0 – 1400 m OHCA based on Argo data (not shown) show that additional cooling occurred between depths of 750 m and 1400 m. As the Argo target sampling depth of 2000 m is achieved by an increasing number of floats, the array will better resolve future deeper changes in OHCA. Variations of pentadal global integrals of OHCA to 3000 m are similar in size and magnitude to annual 0-700 m estimates [*Levitus et al.*, 2005], suggesting that most of the interannual warming and cooling signals are found in the upper 700 m. Still, deepening of the warm bowls in subtropical gyres [*Roemmich et al.*, 2006] and/or the warming of bottom water formed in high latitudes [*Østerhus and Gammelsrød*, 1999; *Johnson and Doney*, 2006] could partially offset the upper ocean cooling. It seems unlikely, however, that the entire signal could be compensated by these processes over such a short period of time.

Assuming that the $3.2 (\pm 1.1) \times 10^{22}$ J was not transported to the deep ocean, previous work suggests that the scale of the heat loss is too large to be stored in any single component of the Earth's climate system [*Levitus et al.*, 2005]. A likely source of the cooling is a small net imbalance in the 340 W/m^2 of radiation that the Earth

exchanges with space. Imbalances in the radiation budget of order 1 W/m^2 have been shown to occur on these time scales and have been related to changes in upper OHCA [Wong *et al.*, 2006]. These findings suggest that the observed decrease in upper ocean heat content from 2003 to 2005 could be the result of a net loss of heat from the Earth to space. Nevertheless, further work will be necessary to determine the exact cause of the cooling.

5. Discussion

This work has several implications. First, the updated time series of ocean heat content presented here (Figure 1) and the newly estimated confidence limits (Figure 3) support the significance of previously reported large interannual variability in globally integrated upper-ocean heat content [Levitus *et al.*, 2005]. However, the physical causes for this type of variability are not yet well understood. Furthermore, this variability is not adequately simulated in the current generation of coupled climate models used to study the impact of anthropogenic influences on climate [Gregory *et al.*, 2004; Barnett *et al.*, 2005; Church *et al.*, 2005; and Hansen *et al.*, 2005]. Although these models do simulate the long-term rates of ocean warming, this lack of interannual variability represents a shortcoming that may complicate detection and attribution of human-induced climate influences.

Changes in OHCA also affect sea level. Sea level rise has a broad range of implications for climate science as well as considerable socioeconomic impacts [Intergovernmental Panel on Climate Change (IPCC), *Climate Change 2001: The Scientific Basis*, 2004]. Diagnosing the causes of past and present sea level change and closure of the sea level budget is therefore a critical component of understanding past changes in sea level as well as projecting future changes. The recent cooling of the upper ocean implies a decrease in the thermosteric component of sea level. Estimates of total sea level [Leuliette *et al.*, 2004; <http://sealevel.colorado.edu>], however, show continued sea-level rise during the past 3 years. This suggests that other contributions to sea-level rise, such as melting of land-bound ice, have accelerated. This inference is consistent with recent estimates of ice mass loss in Antarctica [Velicogna and Wahr, 2006] and accelerating ice mass loss on Greenland [Rignot *et al.*, 2006] but closure of the global sea level budget cannot yet be achieved. New satellite observations from the Gravity Recovery and Climate Experiment (GRACE; launched in March, 2002 and administered by NASA and Deutsches Zentrum für Luft-und Raumfahrt, GRACE will map Earth's gravity field approximately once every 30 days during its lifetime) should soon provide sufficient observations of the redistribution of water mass to more fully describe the causes of recent sea-level change.

Finally, the estimates presented here are made possible only by recent improvements in the global ocean observing system. The sharp decrease in the error since 2002 is due to the dramatic improvement of in situ sampling provided by the Argo array of autonomous profiling CTD floats, and the real-time reporting of Argo data made it possible to extend the estimate through 2005. Characterization of the error budget, which is of paramount importance in the estimate of such globally averaged quantities, was made feasible by the long-term maintenance of high quality altimeter missions such as TOPEX/Poseidon and Jason. The issues relating to sea level rise and the global water budget can only be addressed when the record of satellite gravity measurement from

GRACE achieves adequate duration. GRACE, Argo, and satellite altimetry are core components of the global ocean observing system. Failure to maintain any one of these observing systems would seriously impair our ability to monitor the World Ocean and to unravel its importance to the climate system.

Acknowledgments. Altimeter products used herein were produced by Ssalto/Duacs as part of the Environment and Climate EU Enact project (EVK2-CT2001-00117) and distributed by Aviso, with support from CNES. The bulk of the in situ data used herein were provided through the World Ocean Database 2001 and the Global Temperature-Salinity Profile Program (<http://www.nodc.noaa.gov>). Float data were collected and made freely available by Argo (a pilot program of the Global Ocean Observing System) and contributing national programs (<http://www.argo.net/>). JML and GCJ were supported by the NOAA Climate Program Office and the NOAA Office of Oceanic and Atmospheric Research. The findings and conclusions in this article are those of the authors and do not necessarily represent the views of the National Oceanic and Atmospheric Administration. This research was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. PMEL contribution 2911. JIMAR Contribution 06-359.

Reference List

- Barnett, T. P., D. W. Pierce, K. M. AchutaRao, P. J. Gleckler, B. D. Santer, J. M. Gregory, and W. M. Washington (2005), Penetration of human-induced warming into the World's Oceans, *Science*, *309*, 284-287.
- Church, J. A., N. J. White, and J. M. Arblaster (2005), Significant decadal-scale impact of volcanic eruptions on sea level and ocean heat content, *Nature*, *438*, 74-77 .
- Curry, R. G., and M. S. McCartney (2001), Ocean gyre circulation changes associated with the North Atlantic Oscillation, *J. Phys. Oceanogr.*, *31*, 3374-3400.
- Deser, C., M. A. Alexander, and M. S. Timlin (1999), Evidence for a wind-driven intensification of the Kuroshio Current Extension from the 1970s to the 1980s, *J. Climate*, *12*, 1697-1706.
- Ducet, N., P. Y. Le Traon, and G. Reverdin (2000), Global high-resolution mapping of ocean circulation from the combination of TOPEX/Poseidon and ERS-1 and -2, *J. Geophys. Res.*, *105*, 19,477.
- Gille, S. (2006), Decadal-scale temperature trends in the Southern Hemisphere, manuscript in preparation.
- Gilson, J., D. Roemmich, and B. Cornuelle (1998), Relationship of TOPEX/Poseidon altimetric height to steric height and circulation in the North Pacific, *J. Geophys. Res.*, *103*, 27,947–27,965.
- Gregory, J. M., H. T. Banks, P. A. Stott, J. A. Lowec, M. D. Palmer (2004), Simulated and observed decadal variability in ocean heat content, *Geophys. Res. Lett.*, *31*, L15312, doi:10.1029/2004GL020258.
- Hansen, J., L. Nazarenko, R. Ruedy, M. Sato, J. Willis, A. Del Genio, D. Koch, A. Lacis, K. Lo, S. Menon, T. Novakov, J. Perlwitz, G. Russell, G. A. Schmidt, and N. Tausnev (2005), Earth's energy imbalance: Confirmation and implications, *Science*, *308*, 1431-1435.
- Intergovernmental Panel on Climate Change (IPCC), Climate Change 2001: The Scientific Basis, J. T. Houghton et al., Eds. (Cambridge Univ. Press, New York, 2001).
- Johnson, G. C. and S. C. Doney (2006), Recent western South Atlantic bottom water warming, *Geophys. Res. Lett.*, *33*, L14614, doi:10.1029/2006GL026769.
- Johnson, G. C., M. J. McPhaden, G. D. Rowe, and K. E. McTaggart (2000), Upper equatorial Pacific Ocean current and salinity variability during the 1996 – 1998 El Nino – La Nina cycle. *J. Geophys. Res.*, *105*, 1037-1053.
- Leuliette, E. W., R. S. Nerem, G. T. Mitchum (2004), Calibration of TOPEX/Poseidon and Jason altimeter data to construct a continuous record of mean sea level change, *Marine Geodesy*, *27*, 79-94.
- Levitus, S. J., I. Antonov, and T. P. Boyer (2005), Warming of the world ocean, 1955-2003, *Geophys. Res. Lett.*, *32*, L02604, doi:10.1029/2004GL021592.
- Østerhus, S., and T. Gammelsrød (1999), The Abyss of the Nordic Seas is warming, *J. Climate*, *12*, 3297-3304.
- Pielke, R. A. (2003), Heat storage within the Earth system, *Bull. Amer. Meteor. Soc.*, *84* (3), 331-335.
- Rignot, E. and P. Kanagaratnam (2006), Changes in the velocity structure of the Greenland ice sheet, *Science*, *311*, 986-990.

- Roemmich, D., J. Gilson, R. Davis, P. Sutton, S. Wijffels, and S. Riser (2006), Decadal spin up of the South Pacific subtropical gyre, *J. Phys. Oceanogr.*, in press.
- Willis, J. K., D. Roemmich, and B. Cornuelle (2004), Interannual variability in upper ocean heat content, temperature, and thermosteric expansion on global scales. *J. Geophys. Res.*, *109*, C12036, doi: 10.1029/2003JC002260c.
- Velicogna, I., and J. Wahr (2006), Measurements of time-variable gravity show mass loss in Antarctica, *Science*, *311*, 1754-1756.
- White, W., and C.-K. Tai (1995), Inferring interannual changes in global upper ocean heat storage from TOPEX altimetry, *J. Geophys. Res.*, *100* (C12), 24,943–24,954.
- Wong, T., B. A. Wielicki, R. B. Lee III, G. L. Smith, and K. A. Bush (2006), Re-examination of the observed decadal variability of Earth radiation budget using altitude-corrected ERBE/ERBS nonscanner WFOV data, *J. Climate*, in press.
- Wunsch, C. (1996), *The Ocean Circulation Inverse Problem*, Cambridge University Press, 442 pp.

Figures




Figure 1. Globally averaged annual OHCA [10^{22} J] in the upper 750 m estimated using in situ data alone from 1993 through 2005 (black line) and using in situ data excluding profiling floats (gray line). Error bars (from Figure 3) reflect the standard error estimates discussed in Section 3. Linear trends are computed from a weighted least square fit [Wunsch, 1996] and reflect the OHCA estimate made using all available profile data. Errors for inset linear trend estimates are quoted at the 95% confidence interval.




Figure 2. Map of OHCA change [W m^{-2}] in the upper 750 m from 2003 to 2005.




Figure 3. Standard error for globally averaged OHCA [10^{22} J] of the upper 750 m from 1955 through 2005. This quantity was estimated using satellite altimeter data maps and the historical sampling patterns for in situ profile data in each year as discussed in the text.




Figure 4. Globally averaged ocean temperature change [°C] from 2003 to 2005 versus depth [m]. Thin black lines represent error bounds determined by scaling the uncertainty in heat content using regression coefficients [Willis *et al.*, 2004].

Supplemental Online Material

For each year's sampling pattern, all data values were set equal to one and then objectively mapped using the same technique as was used to produce the OHCA maps. The resulting maps vary between zero and one and illustrate the fraction of variance that can be accounted for using the in situ data in the annual maps of OHCA. The maps for 2002 and 2005 (Figure S1) reveal the reason that the in situ error for the global average OHCA in 2002 is larger than in 2005. In 2002, large swaths of the southern hemisphere and other regions are badly under-sampled. However, by 2005, the growing Argo Project array of profiling CTD floats is doing a much better job at sampling OHCA for the global ice-free ocean [text adapted from *Johnson et al.*, 2006].




Figure S1. Fraction of spatially uniform variance in annual in situ OHCA maps for 2002 (top), a poorly sampled year in terms of in situ data distributions, and 2005 (bottom), a much better sampled year with the advent of the Argo Project (data for 2005 in this plot alone only extend to the end of October).

Supplement Reference List

Johnson, G. C., S. Levitus, J. M. Lyman, C. Schmid, and J. K. Willis (2006), Ocean heat content variability, in *Annual Report on the State of the Ocean and the Ocean Observing System for Climate: FY 2005*, edited by J. M. Levy D. M. Stanitski, and P. Arkin, pp. 74-84, NOAA Climate Program Office, Silver Spring, MD.